


Knots, four dimensions, and fractals

Arunima Ray
Brandeis University


SUNY Geneseo Mathematics Department Colloquium

January 29, 2016


Examples of knots



Examples of knots



Mathematical knots




Take a piece of string, tie a knot in it, glue the two ends together.

Definition

A (mathematical) knot is a closed curve in space with no self-intersections.

Examples of knots



Examples of knots




Figure: Knots in circular DNA.


(Images from Cozzarelli, Sumners, Cozzarelli, respectively.)

The origins of mathematical knot theory

1880's: The æther hypothesis. Lord Kelvin (1824–1907) hypothesized that atoms were 'knotted vortices' in æther.

The origins of mathematical knot theory


1880's: The æther hypothesis. Lord Kelvin (1824–1907) hypothesized that atoms were 'knotted vortices' in æther. This led Peter Tait (1831–1901) to start tabulating knots.



Tait thought he was making a periodic table!

The origins of mathematical knot theory

1880's: The æther hypothesis. Lord Kelvin (1824–1907) hypothesized that atoms were 'knotted vortices' in æther. This led Peter Tait (1831–1901) to start tabulating knots.



Tait thought he was making a periodic table! This view was held for about 20 years (until the Michelson–Morley experiment).

Modern knot theory

Nowadays knot theory is a subset of the field of topology.

Theorem (Lickorish–Wallace, 1960s)

Any 3–dimensional ‘manifold’ can be obtained from \mathbb{R}^3 by performing an operation called ‘surgery’ on a collection of knots.

Modern knot theory

Nowadays knot theory is a subset of the field of topology.

Theorem (Lickorish–Wallace, 1960s)

Any 3–dimensional ‘manifold’ can be obtained from \mathbb{R}^3 by performing an operation called ‘surgery’ on a collection of knots.

Modern knot theory has applications to algebraic geometry, statistical mechanics, DNA topology, quantum computing,

Big questions in knot theory

- ① How can we tell if two knots are equivalent?

Big questions in knot theory

- ① How can we tell if two knots are equivalent?




Figure: These are all pictures of the same knot!

Big questions in knot theory

- ① How can we tell if two knots are equivalent?




Figure: These are all pictures of the same knot!

Big questions in knot theory

- ① How can we tell if two knots are equivalent?




Figure: These are all pictures of the same knot!

- ② How can we tell if two knots are distinct?

Big questions in knot theory

- ① How can we tell if two knots are equivalent?





Figure: These are all pictures of the same knot!

- ② How can we tell if two knots are distinct?
- ③ Can we quantify the 'knottedness' of a knot?

Genus of a knot

Proposition (Frankl–Pontrjagin, Seifert, 1930's)


Any knot bounds a surface in \mathbb{R}^3 .




Genus of a knot

Fundamental theorem in topology


Surfaces are classified by their genus.



genus= 0



genus= 1




genus= 2


Genus of a knot

Fundamental theorem in topology


Surfaces are classified by their genus.



genus= 0



genus= 1



genus= 2

Definition

The genus of a knot K , denoted $g(K)$, is the least genus of surfaces bounded by K .

Genus of a knot

Proposition

If K and J are equivalent knots, then $g(K) = g(J)$.

Genus of a knot

Proposition

If K and J are equivalent knots, then $g(K) = g(J)$.

Proposition

A knot is the unknot if and only if it is the boundary of a disk.

That is, K is the unknot if and only if $g(K) = 0$.

Genus of a knot

Proposition

If K and J are equivalent knots, then $g(K) = g(J)$.

Proposition

A knot is the unknot if and only if it is the boundary of a disk.

That is, K is the unknot if and only if $g(K) = 0$.

If T is the trefoil knot, $g(T) = 1$. Therefore, the trefoil is not equivalent to the unknot.

Connected sum of knots



Figure: The connected sum of two trefoil knots, $T \# T$

Connected sum of knots



Figure: The connected sum of two trefoil knots, $T \# T$

Proposition

Given two knots K and J , $g(K \# J) = g(K) + g(J)$.

Connected sum of knots



Figure: The connected sum of two trefoil knots, $T\#T$

Proposition

Given two knots K and J , $g(K\#J) = g(K) + g(J)$.

Therefore, $g(\underbrace{T\#\cdots\#T}_n) = n$
 n copies

Connected sum of knots



Figure: The connected sum of two trefoil knots, $T\#T$

Proposition

Given two knots K and J , $g(K\#J) = g(K) + g(J)$.

Therefore, $g(\underbrace{T\#\cdots\#T}_n \text{ copies}) = n$

Corollary: There exist infinitely many distinct knots!

Connected sum of knots



Figure: The connected sum of two trefoil knots, $T\#T$

Proposition

Given two knots K and J , $g(K\#J) = g(K) + g(J)$.

Therefore, $g(\underbrace{T\#\cdots\#T}_n) = n$
 n copies

Corollary: There exist infinitely many distinct knots!

Corollary: We can never add together non-trivial knots to get a trivial knot.

Slice knots

Recall that a knot is equivalent to the unknot if and only if it is the boundary of a disk in \mathbb{R}^3 .

Definition

A knot K is slice if it is the boundary of a disk in $\mathbb{R}^3 \times [0, \infty)$.




Figure: Schematic picture of the unknot

Slice knots

Recall that a knot is equivalent to the unknot if and only if it is the boundary of a disk in \mathbb{R}^3 .

Definition

A knot K is slice if it is the boundary of a disk in $\mathbb{R}^3 \times [0, \infty)$.




Figure: Schematic picture of the unknot

Slice knots

Recall that a knot is equivalent to the unknot if and only if it is the boundary of a disk in \mathbb{R}^3 .

Definition

A knot K is slice if it is the boundary of a disk in $\mathbb{R}^3 \times [0, \infty)$.






Figure: Schematic picture of the unknot and a slice knot


Examples of slice knots




Examples of slice knots



Examples of slice knots




Examples of slice knots



Knots of this form are called *ribbon knots*.

Examples of slice knots




Knots of this form are called *ribbon knots*.

Knots, modulo slice knots, form a group called the *knot concordance group*, denoted \mathcal{C} .

Fractals


Fractals are objects that exhibit ‘self-similarity’ at arbitrarily small scales.



i.e. there exist families of injective functions from the set to smaller and smaller subsets (in particular, the functions are non-surjective).

Fractals

Fractals are objects that exhibit ‘self-similarity’ at arbitrarily small scales.



i.e. there exist families of injective functions from the set to smaller and smaller subsets (in particular, the functions are non-surjective).

Conjecture (Cochran–Harvey–Leidy, 2011)

The knot concordance group \mathcal{C} is a fractal.

Satellite operations on knots




Figure: The satellite operation on knots

Satellite operations on knots




Figure: The satellite operation on knots

Any knot P in a solid torus gives a function on the knot concordance group,

$$P : \mathcal{C} \rightarrow \mathcal{C}$$
$$K \mapsto P(K)$$

These functions are called *satellite operators*.

The knot concordance group has fractal properties

Theorem (Cochran–Davis–R., 2012)

Large (infinite) classes of satellite operators $P : \mathcal{C} \rightarrow \mathcal{C}$ are injective.

The knot concordance group has fractal properties

Theorem (Cochran–Davis–R., 2012)

Large (infinite) classes of satellite operators $P : \mathcal{C} \rightarrow \mathcal{C}$ are injective.

Theorem (R., 2013)

There are infinitely many satellite operators P and a large class of knots K such that $P^i(K) \neq P^j(K)$ for all $i \neq j$.

The knot concordance group has fractal properties

Theorem (Cochran–Davis–R., 2012)

Large (infinite) classes of satellite operators $P : \mathcal{C} \rightarrow \mathcal{C}$ are injective.

Theorem (R., 2013)

There are infinitely many satellite operators P and a large class of knots K such that $P^i(K) \neq P^j(K)$ for all $i \neq j$.

Theorem (Davis–R., 2013)

There exist satellite operators that are bijective on \mathcal{C} .

The knot concordance group has fractal properties

Theorem (Cochran–Davis–R., 2012)

Large (infinite) classes of satellite operators $P : \mathcal{C} \rightarrow \mathcal{C}$ are injective.

Theorem (R., 2013)

There are infinitely many satellite operators P and a large class of knots K such that $P^i(K) \neq P^j(K)$ for all $i \neq j$.

Theorem (Davis–R., 2013)

There exist satellite operators that are bijective on \mathcal{C} .

Theorem (A. Levine, 2014)

There exist satellite operators that are injective but not surjective.

Fractals

What is left to show?

In order for \mathcal{C} to be a fractal, we need some notion of distance, to see that we have smaller and smaller embeddings of \mathcal{C} within itself.

That is, we need to exhibit a metric space structure on \mathcal{C} . There are several natural metrics on \mathcal{C} , but we have not yet found one that works well with the current results on satellite operators. The search is on!